

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/92324/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Diz, Paula and Barker, Stephen ORCID: <https://orcid.org/0000-0001-7870-6431>
2016. Approaches and constraints to the reconstruction of palaeoproductivity from Cape Basin abyssal benthic foraminifera (South Atlantic). *Journal of Micropalaeontology* 35 , pp. 195-204. 10.1144/jmpaleo2015-045 file

Publishers page: <http://dx.doi.org/10.1144/jmpaleo2015-045>
<<http://dx.doi.org/10.1144/jmpaleo2015-045>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



**Approaches and constraints to the reconstruction of palaeoproductivity
from Cape Basin abyssal benthic foraminifera (South Atlantic)**

Short title: Palaeoproductivity abyssal benthic foraminifera

Paula Diz¹ and Stephen Barker²

¹Department of Geociencias Marinas y Ordenación del Territorio, Facultad de Ciencias
del Mar, Universidad de Vigo, Campus Lagoas-Marcosende, 36310, Vigo, Spain.

Corresponding author

²School of Earth and Ocean Science, Cardiff University, Cardiff C10 3AT, United
Kingdom

ABSTRACT:

The characteristics of benthic foraminiferal assemblages from well oxygenated deep-sea settings have been used to reconstruct past productivity conditions. None of the different approaches that have been developed are without complications or apply in all settings. In this study we assess the use of benthic foraminifera (accumulation rates and assemblages composition) as proxies for palaeoproductivity changes during the last glacial period (25-95 kyr) in an abyssal core located in the south of Cape Basin (41.1°S, 7.8°E, 4981 m water depth). Assemblage characteristics indicate a generally food-limited environment receiving episodic inputs of labile organic carbon of variable strength. High seasonality in the delivery of organic material to the sea floor in the form of phytodetritus influences the assemblage characteristics because the corresponding response does not involve the whole community. Benefiting from this occasionally high organic input is the opportunistic species, *Epistominella exigua*, that reproduces rapidly to build up large populations. In general, the rest of the species (i.e., less opportunistic compared to *E. exigua*) show only subtle variations in their population densities and fauna composition. Under those circumstances benthic foraminiferal accumulation rates seem to be independent of the amount of organic flux arriving at the sediment surface and respond instead to the strength of phytoplankton blooms.

Key words: benthic foraminifera, palaeoproductivity, seasonality, abyssal, phytodetritus

INTRODUCTION

The fossil record of benthic foraminifera has been widely used as a proxy for past environmental conditions (for reviews see Gooday 2003; Gooday & Jorissen 2012; Jorissen *et al.* 2007; Murray 2006). Despite the complexity of factors playing a role in foraminiferal distribution and the effect of post-depositional processes (Jorissen *et al.* 2007; Murray 2001), there is a general agreement that benthic foraminifera from deep sea environments largely respond to a few parameters: the organic carbon flux, bottom-water hydrodynamics (and related grain size), bottom and pore-water oxygenation and carbonate saturation (see reviews in Mackensen *et al.*, 1995; Jorissen *et al.* 2007). Within these parameters, the estimation of past changes in the primary production (and organic carbon flux to the sea floor) from benthic foraminifera has received major attention in palaeoceanographic studies because of the implications of changes in the biological pump on past and future climate. Based on knowledge of the ecology of benthic foraminifera, a number of studies have suggested the abundance of benthic foraminifera (benthic foraminiferal accumulation rates, Herguera & Berger 1991), the contribution of different foraminiferal morphotypes (Corliss & Chen 1988) and species composition (e.g., Altenbach *et al.* 1999) as quantitative and qualitative proxies for past organic carbon fluxes arriving at the seafloor. Faunal composition might also provide information about the quality (refractory versus labile) of the organic carbon (e.g., Fontanier *et al.* 2005). The input of labile organic carbon to the sea floor derived from seasonal or episodic phytoplankton blooms at the surface also influences assemblage composition (e.g., Smart *et al.* 1994; Sun *et al.* 2006) and diversity (e.g., Corliss *et al.* 2009; Enge *et al.* 2011; Gooday *et al.* 2010, 2012) so providing information about the periodicity of the organic carbon flux.

Evaluating the extent to which benthic foraminifera can be used to infer a particular set of past environmental conditions and understanding the complicating factors and limitations of such reconstructions is fundamental to advancing the use of proxy methods based on these

organisms. In this study we discuss palaeoenvironmental inferences based on benthic foraminiferal assemblage characteristics from an abyssal core located in the south of Cape Basin (South Atlantic). We examine the consistency of the most typical palaeoproductivity proxies by comparing information provided by our assemblages with environmental parameters from independent proxies obtained in the same core and other proxy archives.

MATERIAL AND METHODS

The marine sediment core discussed in this study (TNO57-21) was recovered in the south Cape Basin, slightly north of the Agulhas ridge in the southeast Atlantic Ocean (41.1°S, 7.8°E, 4981 m of water depth, 13.8 m long, Fig. 1) . It is presently bathed by the poorly ventilated, cold and fresh Antarctic Bottom Water (AABW). The location lies below at the Subtropical Convergence (STC) front that separates the cold low-salinity Subantarctic waters to the south and warm saline Subtropical waters to the north. The complex interaction of physico-chemical factors associated with cross-frontal mixing of subantarctic and subtropical waters and the intense turbulence and eddy activity (Baker-Yeboah *et al.* 2010; Llido *et al.* 2004) results in relatively elevated pulses or episodes of primary production in the surface waters of the study area (Froneman *et al.* 1997; Machu *et al.* 2005).

Several previous micropalaeontological studies have been carried out on core TN057-21. The palaeoceanographic significance of planktonic foraminifera assemblages are discussed in Barker *et al.* (2009, 2010) and (Barker & Diz 2014). The linkages between the record of the benthic foraminifera *Epistominella exigua* and millennial-scale climate variability during the last 95 kyr are considered in Diz & Barker (2015). However, detailed information about benthic assemblage composition from ~25 to 95 kyr, and relevant additional palaeoenvironmental information, were beyond the scope of these earlier papers and are therefore addressed in the present study. The age model used in this study is described in

Barker & Diz (2014) and the methods for the study of benthic fauna are indicated in Diz & Barker (2015). Briefly, foraminifera are picked from the > 125 µm fraction, mounted on faunal slides and identified following the generic assignments of (Loeblich & Tappan 1987) and published illustrations (see Table 1). The study of benthic foraminifera in the >125 µm size fraction potentially underestimate the abundance of small (63-125 µm) opportunistic taxa such as *Alabaminella weddellensis* which are typically abundant in areas characterized by episodic phytoplankton blooms (e.g., Diz *et al.* 2007; Smart *et al.* 2010; Thomas *et al.* 1995). On the other hand, it allows for studying a feasible number of samples still showing major faunal patterns (e.g., Poli *et al.* 2012, Schönfeld *et al.* 2012). Raw counts can be found in Diz & Barker (2015) and the most characteristic species of benthic foraminifera are illustrated in this study (Plate I). The assemblages are largely dominated by *Epistominella exigua* (>40% on average), a phytodetritus related species. In order to assess the ecological significance of the other relevant species we discuss the assemblages after removing the contribution of *E. exigua* (i.e., the "phytodetritus free assemblage", see also Thomas *et al.* 1995). Additionally, we examined the record of benthic foraminiferal accumulation rate (BFAR, total number of foraminifera x cm⁻² x kyr⁻¹) and phytodetritus free benthic foraminiferal accumulation rate (BFAR_{phfree}). BFARs are calculated as the number of individuals per gram of total dry bulk sediment (#bf x g⁻¹, Diz & Barker 2015) x linear sedimentation rates (cm x kyr⁻¹) x density (g x cm⁻³). Sedimentation rates are taken from the age model developed by Barker & Diz (2014) and vary from 8 to 16 cm kyr⁻¹. It is considered that the age model is sufficiently robust as to assume that differences in BFAR are not mainly caused by inaccuracies in the calculation of sedimentation rates. In the absence of direct measurements of dry or wet bulk density, we used the algorithm developed by Sachs & Anderson (2003) for the calculation of, which is based on the percentage of calcium carbonate in order to calculate densities in core TNO57-21. To do this, the high resolution % CaCO₃ record of core TNO57-21 (Sachs & Anderson

2003) was re-sampled and values interpolated to the lower resolution record of benthic foraminiferal abundance using the program Analyseries (Paillard *et al.* 1996). Then, density values were calculated using Sachs and Anderson' formula. Calculated values vary between 0.5 to 0.7 g x cm⁻³.

The core TNO57-21 was recovered from deep waters that are currently undersaturated with respect to calcium carbonate (4891 m water depth, see Supplementary Information by Barker *et al.* 2010). Several episodes of increased carbonate dissolution have been identified in planktonic foraminiferal (Barker & Diz 2014; Barker *et al.* 2009, 2010) as well as benthic foraminiferal (Diz & Barker 2015) assemblages. By considering the information provided by multiple proxies (e.g., % of coarse fraction, % CaCO₃, foraminiferal abundances, composition of assemblages) it was indicated that carbonate dissolution does not seem to exert an overall control on benthic abundance or species distribution with the exception of the Holocene (not considered in this study) and Greenland Stadial (GS) events GS19 and 21. Those poorly preserved periods (indicated in figure 2 by bars) are characterized by low percent of calcium carbonate, very low foraminiferal abundances and low number of benthic taxa. They are not used to infer palaeoenvironmental conditions (see detailed discussion in Diz & Barker 2015). The comparison of the composition of the "dissolved assemblages" with modern samples (Mackensen *et al.* 1993) indicates that *Nutallides umboniferus* is very rare along core TNO57-21 (see Plate). *N. umboniferus* is generally related to corrosive bottom waters in the South Atlantic (Mackensen *et al.* 1993; Harloff & Mackensen, 1997; Schmied *et al.* 1997). It might suggest that other factors more than carbonate undersaturated waters are playing a role in this species distribution (i.e, oligotrophic conditions and absence of phytodetritus deposition, Smart & Gooday, 1997; Kurbejewit *et al.* 2000; Carman & Keigwin, 2004).

RESULTS

The foraminifera fauna of core TNO57-21 comprises 39 benthic species dominated by calcareous forms. The Shannon H diversity Index averages 1.9 (Figure 2a). High diversity values occur when the relative contribution of the dominant phytodetritus-related species *Epistominella exigua* (Brady) are diminished. The averaged relative contribution of this species is as high as 47% (Fig. 2b). Other, relatively common species are *Pullenia osloensis* (Feyling-Hanssen) (average percentage 11.9%, Fig. 2d), *Oridorsalis umbonatus* (Reuss) (6%, Fig. 2h), *Fontbotia wuellerstorfi* (Schwager) (7.9%, Fig. 2g) and the group of species belonging to the suborder *Lagenina* called "Lagenina group" (see Table 1, 11.9%, Fig. 2c). Secondary calcareous species (<10 % on average) showing high abundances at particular intervals are *Pyrgo murrinha* (Schwager) (Fig. 2j), *Globocassidulina subglobosa* (Brady) (Fig. 2i), *Melonis* spp. (comprised of *M. zaandamae* (Van Voothuysen), *Melonis pompilioides* (Fichtel and Moll) and *Melonis* sp., Fig. 2f) and *Rutherfordoides tenuis* (Phleger and Parker) (Fig. 2l). Calcareous species with short-lived peaks are *Uvigerina auberiana* (d'Orbigny) (Fig. 2m) and *Quinqueloculina* cf. *seminula* (Fig. 2n). The most characteristic agglutinated foraminifera is the calcareous agglutinated *Siphotextularia rolshauseni* (Phleger and Parker, Fig. 2e) with *Eggerella bradyi* (Cushman) as secondary species (Fig. 2k).

The BFAR values in core TNO57-21 show large fluctuations ranging from 30 to 1100 foraminifera ($> 125 \mu\text{m}$ fraction) $\text{cm}^{-2} \text{kyr}^{-1}$. On average, peak BFAR values are higher during northern stadial events occurring during MIS 3 than over MIS 5 (Fig. 3a). The lowest BFAR values occur during GS19 and GS21 (also the Holocene, not shown) when benthic foraminiferal dissolution was inferred (Diz & Barker 2015) and thus they are excluded from interpretation (Fig. 3a). In general, episodes of peak BFAR values are largely driven by the phytodetritus-related species *Epistominella exigua* (Fig. 3a), and show a tight temporal correspondence with abrupt climate changes occurring in the North Atlantic Ocean and concomitant antiphase response in the South Atlantic (see Fig. 3 and Diz & Barker 2015).

Episodes of high $\text{BFAR}_{\text{phfree}}$ coincide with BFAR although they are of much lower magnitude and sharpness (Fig. 3a, grey shaded area). The highest $\text{BFAR}_{\text{phfree}}$ values coincide with a single peak in *Quinqueloculina cf. seminula* (Fig. 2n) at ~45 kyr and near the beginning of MIS 3 (~56-58 kyr) with high relative abundances of *G. subglobosa*, *F. wuellerstorfi* and *O. umbonatus* (see Figure 2).

DISCUSSION

Reconstructing the flux of organic carbon to the seafloor from benthic foraminifera

Benthic foraminiferal accumulation rates. The benthic foraminiferal accumulation rate (BFAR, number of foraminifera $\times \text{cm}^{-2} \times \text{kyr}^{-1}$) has been suggested as a palaeoproductivity proxy (Herguera 1992, 2000; Herguera & Berger 1991). However, several studies indicated that obtaining a quantitative estimation of primary production or export production from BFAR values might be limited by several factors, with the difficulty of obtaining reliable calibrations, carbonate dissolution and taphonomical processes being the most important (see review in Jorissen *et al.* 2007). In this study no attempts have been made to estimate absolute palaeoproductivity or palaeo-carbon fluxes, but we evaluate the potential use of the calculated BFAR as a semi-quantitative proxy for food supply to the ocean floor (e.g., Thomas *et al.* 1995; Alegret & Thomas 2009) in core TNO57-21.

The BFAR values during the Holocene and late deglaciation (not plotted but calculated on the basis of Diz & Barker (2015) data; <100 foraminifera $\text{cm}^{-2} \text{kyr}^{-1}$) are within the range of values for abyssal core tops from the Pacific open ocean at water depths > 4000 m likely affected by carbonate dissolution (Herguera 1992; Herguera & Berger 1991). However, peak BFAR values from MIS 3 to MIS 5 are four to eight times Holocene values (Figure 3). These accumulation rates are substantially higher than maxima values recorded in core-top samples

from widely distributed open-ocean areas at various depths in the Pacific. Following the use of BFAR as a palaeoproductivity proxy, these results might suggest past increases in the organic carbon flux to the sea floor up to several times the present conditions. According to modern primary production distributions (see Machu *et al.* 2005, their Fig. 5) this might be equivalent to a change from oligotrophic subtropical open ocean areas to high productivity upwelling centres (e.g., Benguela upwelling). In fact, peaks of BFAR reach values found in continental shelf areas affected by seasonal upwelling such as Cape Blanc upwelling (Guichard *et al.* 1999; Zarriess & Mackensen 2010) or the Southwest African upwelling (Schmiedl & Mackensen 1997). Such a change in the palaeoenvironmental conditions, if it had occurred, would have involved a large change in assemblage composition, i.e. from low productivity faunas (low abundances) to a high productivity assemblage (high BFAR). This seems to be unrealistic considering that assemblage changes are subtle in comparison to BFAR fluctuations (see Figure 2-3) and they do not involve the appearance of species related to eutrophic conditions (see next section). In fact, variations in BFAR are mainly driven by the phytodetritus-related species *Epistominella exigua* (Table 1, Figure 3a). In the absence of this species fluctuations in BFAR_{phfree} (BFAR phytodetritus free; "phfree") are of much lower magnitude and suggest only subtle changes in organic carbon fluxes.

This could be explained by the fact of that when large quantities of fresh marine organic matter (phytodetritus) are available, even for short periods of time, opportunistic species, such as *Epistominella exigua*, reproduce rapidly and produce a large number of tests (see Thiel *et al.* 1989). The accumulation of foraminifera is not related in a predictable manner to organic flux and, in these circumstances, BFAR should not be used as a quantitative (or semiquantitative) proxy (Thomas *et al.*, 1995; Diz *et al.*, 2007; Smart, 2008; Zarriess & Mackensen, 2010) for the organic carbon flux reaching the sea floor. BFAR is rather related to the strength of the phytoplankton blooms. The episodic and labile nature of the

phytodetritus aggregates reaching the sea floor (Beaulieu 2002; Smith *et al.*, 2008) could explain the weak correlation ($r = -0.24$, $p = 0.000$, $n = 467$) between BFAR (also $\text{BFAR}_{\text{phfree}}$) and the alkenones flux, a proxy for averaged organic carbon flux to the seafloor (Sachs & Anderson 2003) obtained in the same core (Fig. 3c).

Benthic foraminiferal assemblage composition. The use of benthic foraminiferal assemblages as palaeo-productivity proxies is based on the relation of faunal composition to organic flux rates (e.g., Morigi *et al.* 2001). Again, the quantification of flux regimes from the relative abundance of major species is complicated by ecological factors, broad adaptability of some species to organic carbon fluxes and uncertainties related to estimations of carbon flux (Altenbach *et al.* 1999; Gooday 2003; Jorissen *et al.* 2007). Thus, the information provided by the faunal composition of core TNO57-21 is used here as qualitative proxy for the organic carbon flux or its quality/periodicity.

The relative abundance of particular species or group of species is related qualitatively to the predominance of a particular flux regime (i.e., high, intermediate, low-flux species), quality (i.e., labile, refractory) or seasonality of the flux (i.e., seasonal, sustained). In this study, groups are defined based on representative species with well-known ecology (see the ecological attributions of characteristic species in Table 1). Because organic flux dependent patterns should be best analysed within the environmental optimum of species (Altenbach *et al.* 1999) those species with percentages lower than 10% for most of the record (e.g., *Eggerella bradyi*, *Quinqueloculina cf. seminula*, *Rutherfordoides tenuis*, see Plate) and/or those species for whom the ecology is not well-constrained are not included in any of the groups. An example of the later is the high contribution to the assemblage of individuals belonging to the "Lagenina group" (>10%, Fig. 2c, see Plate). Unfortunately their ecology is

poorly known (see Table 1) and consequently the contribution to the paleoenvironmental understanding of the assemblage changes is limited.

The most abundant species throughout the core is *Epistominella exigua*, an epifaunal/shallow infaunal abyssal deep-sea opportunistic foraminifera that flourishes and rapidly reproduces in the presence of phytodetritus aggregates (see ecological attributions in Table 1). Thus, high relative abundance of *E. exigua* (Figure 2b) along core TNO57-21 indicates overall low organic carbon fluxes punctuated by episodic phytoplankton blooms that resulted in the deposition of phytodetrital inputs (i.e., labile organic carbon). This interpretation is supported by the "phytodetritus free" assemblage which is composed by species typical of oligotrophic ("low flux") to moderately mesotrophic ("intermediate") environmental conditions.

The "low-flux species" are the epifaunal *Fontbotia wuellerstorfi*, *Pyrgo murrinha* and the shallow infaunal *Oridorsalis umbonatus* and *Globocassidulina subglobosa* (see Table 1). All together these represent > 30% of the phytodetritus free assemblage (Fig. 3f) reaching up to 60% during early MIS3. For some of these species (i.e. *G. subglobosa*, *F. wuellerstorfi*, *P. murrinha*) a relationship to phytodetritus input (See Table 1) has also been suspected.

However, their downcore relative abundance does not covary with BFAR or percentages of phytodetritus-related species *E. exigua* (Figure 3). Instead, they must respond to a different type of phytodetritus or, more likely, they prefer sustained (and low) organic carbon flux to the sea floor rather than pulsed (see also Smart, 2008).

The group of species considered indicative of an "intermediate -flux" regime are mainly represented by the relatively common shallow infaunal *Pullenia osloensis* together with *Melonis* spp. and *Siphotextularia rolshauseni* (Figure 3e, Table 1). Those species show a statistically significant negative correlation ($p < 0.000$) to the relative contribution of the species belonging to low - flux group ($r_{O. umbon-S. rolshauseni} = -0.38$; $r_{P. osloensis-P. murrinha} = -0.39$; $r_{P. osloensis-F. wuellerstorfi} = -0.28$; $r_{P. osloensis-G. subglobosa} = -0.23$; $r_{O. umbonatus-Melonis spp.} = -0.35$). The relative

contribution of the "intermediate-flux" group indicates slightly more eutrophic conditions during MIS 5 and MIS 4 than during MIS 3. Peaks of the intermediate flux fauna occurring over MIS 4 are mainly caused by noticeably increased contributions of *Melonis* spp. and *S. rolshauseni* (Figure 2d-f, Figure 3e) occurring in coincidence with high relative contribution of *Epistominella exigua* (Fig. 2-3) and the substantial increase in the alkenone flux measured during glacial MIS 4 (Fig. 3c). These two species show affinity for degraded organic carbon (Table 1) suggesting that during MIS 4 part of the sedimentary organic compounds are of low nutritional value. This might be explained by a different type and/or fate of phytodetrital material arriving to the seafloor which could promote the response of species that benefit from the bloom indirectly (i.e., bacteria colonizing aggregates, Koho *et al.*, 2008). In fact, it has been inferred (Diz & Barker 2015) that a slightly different physical conditions promoting phytoplankton blooms in the surface waters occurred during MIS 4/MIS 5 transition and early MIS 4 (seasonality and windiness) than during MIS 3 and late MIS 5 (mainly eddy activity). Notably, that assemblage change does not trigger a parallel increase in the $\text{BFAR}_{\text{phfree}}$ indicating that a change in the food quality does not force a change in the absolute numbers of benthic foraminiferal accumulation rate but only in the assemblage composition.

CONSTRAINTS AND APPROACHES: OVERVIEW

Benthic foraminifera (BFAR and assemblage composition) were analysed in abyssal core TN057-21. Episodic phytoplankton blooms represents the main source of "disturbance" to the benthic environment and have important consequences for the structure and taxonomic composition of the benthic foraminifera record. The inferred phytodetritus deposition does not

seem to trigger a response by the whole community but it is limited to a single species, leading to a decrease in the diversity of the assemblages. The species benefiting from phytodetritus input is the opportunistic (r-strategist) *Epistominella exigua* which reacts quickly, producing high numbers of individuals and consequently high benthic foraminiferal accumulation rates. The response of the remaining fauna (the so called "phytodetritus free assemblage") to phytodetritus input is mainly constrained to a concomitant moderate increase in the foraminiferal abundance ($\text{BFAR}_{\text{phfree}}$) with overall little variations in the assemblage composition. This could be explained by the different ecological preferences of species with lower reproductive potential (k-strategists) and likely benefiting from other type organic carbon (less labile). All these data together suggest BFAR could not be a reliable proxy for palaeoproductivity in abyssal (food limited) environments when benthic foraminifera are highly dependent on the rate and nature of the input of labile organic material generated in the euphotic zone. Under these circumstances fluctuations in BFAR do not reflect the flux of organic carbon to the seafloor but rather the strength of phytoplankton blooms in the surface waters. In this case, the assemblage composition is relevant for interpreting palaeoenvironmental conditions.

ACKNOWLEDGEMENTS

Authors wish to thank the reviewers Christopher Smart and Andrew Gooday for their suggestions that improved the manuscript. This work was supported by the UK Natural Environmental Research Council NE/J008133/1 and R+D contract CO-078-10 (Ref. UVigo). P. D. was supported by Xunta de Galicia (Programa Investigadores Isidro Parga Pondal) and University of Vigo (Programa Retención de Talento 2015).

323

324 **REFERENCES**

325 Alegret, L. & Thomas, E. 2009. Food supply to the seafloor in the Pacific Ocean after the
326 Cretaceous/Paleogene boundary event. *Marine Micropaleontology*, **73**(1-2): 105-116

327 Altenbach, A.V., Pflaumann, U., Schiebel, R., Thies, A., Timm, S. & Trauth, M. 1999.
328 Scaling percentages and distributional patterns of benthic foraminifera with flux rates of
329 organic carbon. *Journal of Foraminiferal Research*, **29**: 173-185.

330 Alve, E. 2010. Benthic foraminiferal responses to absence of fresh phytodetritus: A two-
331 year experiment. *Marine Micropaleontology*, **76**(3-4): 67-75.

332 Anderson, J.B. 1975. Factors controlling CaCO₃ dissolution in the Weddell sea from
333 foraminiferal distribution patterns. *Marine Geology*, **19**: 315-332.

334 Baker-Yeboah, S., Byrne, D.A. & Watts, D.R. 2010. Observations of mesoscale eddies in
335 the South Atlantic Cape Basin: Baroclinic and deep barotropic eddy variability. *Journal of*
336 *Geophysical Research: Oceans*, **115**(C12): C12069, doi: 10.1029/2010jc006236.

337 Barker, S., Diz, P., Vautravers, M.J., Pike, J., Knorr, G., Hall, I.R. & Broecker, W.S.
338 2009. Interhemispheric Atlantic seesaw response during the last deglaciation. *Nature*,
339 **457**(7233): 1097-1102.

340 Barker, S., Knorr, G., Vautravers, M.J., Diz, P. & Skinner, L.C. 2010. Extreme deepening
341 of the Atlantic overturning circulation during deglaciation. *Nature Geoscience*, **3**(8): 567-571.

342 Barker, S., Knorr, G., Edwards, R.L., Parrenin, F., Putnam, A.E., Skinner, L.C., Wolff, E.
343 & Ziegler, M. 2011. 800,000 Years of Abrupt Climate Variability. *Science*, **334**(6054): 347-
344 351.

345 Barker, S. & Diz, P. 2014. Timing of the descent into the last Ice Age determined by the
346 bipolar seesaw. *Paleoceanography*: 2014PA002623, **29**(6): 489-507.

347 Beaulieu, S.E. 2002. Accumulation and fate of phytodetritus on the sea floor.
348 *Oceanography and Marine Biology Annual Review*, **40**: 171-232.

349 Boltovskoy, E. & Boltovskoy, D. 1989. Paleocene-Pleistocene Benthic Foraminiferal
350 Evidence of Major Paleoceanographic Events in the eastern South Atlantic (DSDP Site 525,
351 Walvis Ridge). *Marine Micropaleontology*, **14**: 283-316.

- 352 Burke, S.K., Berger, W.H., Coulbourn, W.T. & Vincent, E. 1993. Benthic foraminifera in
353 box core ERDC 112, Ontong Java Plateau. *Journal of Foraminiferal Research*, **23**(1): 19-39,
354 doi: 10.2113/gsjfr.23.1.19.
- 355 Caralp, H.M. 1989. Abundance of *Bulimina exilis* and *Melonis barleeanum*: relationship
356 to the quality of marine organic matter. *Geo-Marine Letters*, **9**: 37-43.
- 357 Carman, M.R. & Keigwin, L.D. 2004. Preservation and color differences in *Nuttallides*
358 *umbonifera* *Journal of Foraminiferal Research*, **34**(2): 102-108.
- 359 Corliss, B.H. & Chen, C. 1988. Morphotype patterns of Norwegian Sea deep-sea benthic
360 foraminifera and ecological implications. *Geology*, **16**: 716-719.
- 361 Corliss, B.H. 1979. Recent deep-sea benthonic foraminiferal distributions in the
362 southeast Indian Ocean: Inferred bottom-water routes and ecological implications. *Marine*
363 *Geology*, **31**(1-2): 115-138.
- 364 Corliss, B.H. 1983. Distribution of Holocene deep-sea benthonic foraminifera in the
365 southwest Indian Ocean. *Deep Sea Research Part A, Oceanographic Research Papers*, **30**(2):
366 95-117.
- 367 Corliss, B.H., Brown, C.W., Sun, X. & Showers, W.J. 2009. Deep-sea benthic diversity
368 linked to seasonality of pelagic productivity. *Deep Sea Research Part I: Oceanographic*
369 *Research Papers*, **56**(5): 835-841.
- 370 Cornelius, N. & Gooday, A.J. 2004. 'Live' (stained) deep-sea benthic foraminiferans in
371 the western Weddell Sea: trends in abundance, diversity and taxonomic composition along a
372 depth transect. *Deep Sea Research Part II: Topical Studies in Oceanography*, **51**(14-16):
373 1571-1602.
- 374 De, S. & Gupta, A.K. 2010. Deep-sea faunal provinces and their inferred environments in
375 the Indian Ocean based on distribution of Recent benthic foraminifera. *Palaeogeography*,
376 *Palaeoclimatology, Palaeoecology*, **291**(3-4): 429-442.
- 377 Diz, P. & Barker, S. 2015. Linkages between rapid climate variability and deep-sea
378 benthic foraminifera in the deep Subantarctic South Atlantic during the last 95 kyr.
379 *Paleoceanography*, **30**(6): 601-611.
- 380 Diz, P., Hall, I.R., Zahn, R. & Molyneux, E.G. 2007. Paleooceanography of the southern
381 Agulhas Plateau during the last 150 ka: Inferences from benthic foraminiferal assemblages
382 and multispecies epifaunal carbon isotopes. *Paleoceanography*, **22**(4): PA4218.

- Eberwein, A. & Mackensen, A. 2006. Regional primary productivity differences off Morocco (NW Africa) recorded by modern benthic foraminiferal and their stable carbon isotopic composition. *Deep-Sea Research I*, **53**: 1379-1405.
- Enge, A.J., Nomaki, H., Ogawa, N.O., Witte, U., Moeseneder, M.M., Lavik, G., Ohkouchi, N., Kitazato, H., Kucera, M. & Heinz, P. 2011. Response of the benthic foraminiferal community to a simulated short-term phytodetritus pulse in the abyssal North Pacific. *Marine Ecology Progress Series*, **438**: 129-142.
- Epica, Community & Members. 2006. One-to-one coupling of glacial climate variability in Greenland and Antarctica. *Nature*, **444**: 195-198.
- Fontanier, C., Jorissen, F.J., Chaillou, G., Anschutz, P., Grémare, A. & Griveaud, C. 2005. Live foraminiferal faunas from a 2800 m deep lower canyon station from the Bay of Biscay: Faunal response to focusing of refractory organic matter. *Deep-Sea Research I*, **52**: 1189-1227.
- Fontanier, C., Jorissen, F.J., Licari, L., Alexandre, A., Anschutz, P. & Carbonel, P. 2002. Live benthic foraminiferal faunas from the Bay of Biscay: faunal density, composition, and microhabitats. *Deep-Sea Research I*, **49**: 751-785.
- Froneman, P.W., Perissinotto, R. & Pakhomov, E.A. 1997. Biogeographical structure of the microphytoplankton assemblages in the region of the Subtropical Convergence and across a warm-core eddy during austral winter. *Journal of Plankton Research*, **19**(4): 519-531.
- Gooday, A.J. 1988. A response by benthic Foraminifera to the deposition of phytodetritus in the deep sea. *Nature*, **332**: 70-73.
- Gooday, A.J. 1993. Deep-sea benthic foraminiferal species which exploit phytodetritus: Characteristic features and controls on distribution. *Marine Micropaleontology*, **22**: 187-205.
- Gooday, A.J. 2003. Benthic foraminifera (protista) as tools in deep-water palaeoceanography: Environmental influences on faunal characteristics. *Advances in Marine Biology*. Academic Press: 1-90.
- Gooday, A.J., Malzone, M.G., Bett, B.J. & Lamont, P.A. 2010. Decadal-scale changes in shallow-infaunal foraminiferal assemblages at the Porcupine Abyssal Plain, NE Atlantic. *Deep Sea Research Part II: Topical Studies in Oceanography*, **57**(15): 1362-1382.
- Gooday, A., Bett, B., Jones, D. & Kitazato, H. 2012. The influence of productivity on abyssal foraminiferal biodiversity. *Marine Biodiversity*, **42**(4) : 1-17.

- Gooday, A.J. & Jorissen, F.J. 2012. Benthic Foraminiferal Biogeography: Controls on Global Distribution Patterns in Deep-Water Settings. *Annual Review of Marine Science*, **4**(1): 237-262.
- Guichard, S., Jorissen, F. & Peypouquet, J.P. 1999. Late Quaternary benthic foraminiferal records testifying lateral variability of the Cape Blanc upwelling signal. *Comptes Rendus de l'Academie de Sciences - Serie Ila: Sciences de la Terre et des Planetes*, **329**(4): 295-301.
- Gupta, A.K. 1997. Paleooceanographic and paleoclimatic history of the Somali Basin during the Pliocene-Pleistocene; multivariate analyses of benthic foraminifera from DSDP Site 241 (Leg 25). *Journal of Foraminiferal Research*, **27**(3): 196-208.
- Gupta, A.K. & Thomas, E. 2003. Initiation of Northern Hemisphere glaciation and strengthening of the northeast Indian monsoon: Ocean drilling program site 758, eastern equatorial Indian Ocean. *Geology*, **31**(1): 47-50.
- Harloff, J. & Mackensen, A. 1997. Recent benthic foraminifera associations and ecology of the Scotia Sea and Argentine Basin. *Marine Micropaleontology*, **31**: 1-29.
- Herguera, J.C. & Berger, W.H. 1991. Paleoproductivity from benthic foraminifera abundance: Glacial to postglacial change in the west-equatorial Pacific. *Geology*, **19**: 1173-1176.
- Herguera, J.C. 1992. Deep-Sea benthic foraminifera and biogenic opal: Glacial to postglacial productivity changes in the western equatorial Pacific. *Marine Micropaleontology*, **19**: 79-98.
- Herguera, J.C. 2000. Last glacial paleoproductivity patterns in the eastern equatorial Pacific: benthic foraminifera records. *Marine Micropaleontology*, **40**: 259-275.
- Jorissen, F.J., Fontanier, C. & Thomas, E. 2007. Chapter Seven Paleooceanographical Proxies Based on Deep-Sea Benthic Foraminiferal Assemblage Characteristics. *Developments in marine geology 1*, Elsevier, 263-325.
- Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster, B., Nouet, J., Barnola, J.M., Chappellaz, J., Fischer, H., Gallet, J.C., Johnsen, S., Leuenberger, M., Loulergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud, D., Schilt, A., Schwander, J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J.P., Stenni, B., Stocker, T.F., Tison, J.L., Werner, M. & Wolff, E.W. 2007. Orbital and

- 444 Millennial Antarctic Climate Variability over the Past 800,000 Years. *Science*, **317**(5839):
445 793-796.
- 446 Koho, K.A., Langezaal, A.M., van Lith, Y.A., Duijnste, I.A.P. & van der Zwaan, G.J.
447 2008. The influence of a simulated diatom bloom on deep-sea benthic foraminifera and the
448 activity of bacteria: A mesocosm study. *Deep-Sea Research I*, **55**: 696-719.
- 449 Kurbjeweit, F., Schimiedl, G., Schiebel, R., Hemleben, C., Pfannkuche, O., Wallmann,
450 K. & Schäfer, P. 2000. Distribution, biomass and diversity of benthic foraminifera in relation
451 to sediment geochemistry in the Arabian Sea. *Deep-Sea Research II*, **47**: 2913-2955.
- 452 Kurihara, K. & Kennett, J.P. 1986. Neogene benthic foraminifera: distribution in depth
453 traverse, southwest Pacific. In: Kennett, J.P., von der Borch, C. C., and others (Ed) Initial
454 Reports of the Deep Sea Drilling Project, Volume 90 (Part 2). US. Government Printing
455 Office, Washington, 1037-1077.
- 456 Linke, P. & Lutze, G.F. 1993. Microhabitat preferences of benthic foraminifera a static
457 concept or a dynamic adaptation to optimize food acquisition? *Marine Micropaleontology*,
458 **20**: 215-234.
- 459 Llido, J., Machu, E., Sudre, J., Dadou, I. & Garçon, V. 2004. Variability of the biological
460 front south of Africa from SeaWiFS and a coupled physical-biological model. *Journal of*
461 *Marine Research*, **62**: 595- 609.
- 462 Lobeguer, M.K. & Sen Gupta, B.K. 2008. Foraminifera of Hydrocarbon seeps, Gulf of
463 Mexico. *Journal of Foraminiferal Research*, **38**(2): 93-116.
- 464 Loeblich, A.R. & Tappan, H. 1987. Foraminifera Genera and Their Classification. Van
465 Nostrand Reinhold, New York.
- 466 Machu, E., Biastoch, A., Oeschies, A., Kawamiya, M., Lutjeharms, J.R.E. & Garçon, V.
467 2005. Phytoplankton distribution in the Agulhas system from a coupled physical-biological
468 model. *Deep-Sea Research Part I: Oceanographic Research Papers*, **52**(7): 1300-1318.
- 469 Mackensen, A., Grobe, H., Kuhn, G. & Fütterer, D.K. 1990. Benthic foraminiferal
470 assemblages from the eastern Weddell Sea between 68 and 73°S: Distribution, ecology and
471 fossilization potential. *Marine Micropaleontology*, **16**: 241-283.
- 472 Mackensen, A., Fütterer, D.K., Grobe, H. & Schimiedl, G.E.P.d. 1993. Benthic
473 foraminiferal assemblages from the eastern South Atlantic Polar Front region between 35° and
474 57°S: Distribution, ecology and fossilization potential. *Marine Micropaleontology*, **22**: 33-69.

- 475 Mackensen, A., Schmiedl, G., Harloff, J. & Giese, M. 1995. Deep-sea foraminifera in the
476 South Atlantic Ocean: Ecology and assemblage composition. *Micropaleontology*, **41**: 342-
477 358.
- 478 Miao, Q. & Thunell, R.C. 1993. Recent deep-sea benthic foraminiferal distributions in
479 the South China and Sulu Seas. *Marine Micropaleontology*, **22**(1-2): 1-32.
- 480 Morigi, C., Jorissen, F.J., Gervais, A., Guishard, S. & Borsetti, A.M. 2001. Benthic
481 foraminiferal faunas in surface sediments off NW Africa: Relationship with organic flux to
482 the ocean floor. *Journal of Foraminiferal Research*, **31**: 350-368.
- 483 Murgese, D.S. & De Deckker, P. 2007. The Late Quaternary evolution of water masses in
484 the eastern Indian Ocean between Australia and Indonesia, based on benthic foraminifera
485 faunal and carbon isotopes analyses. *Palaeogeography, Palaeoclimatology, Palaeoecology*,
486 **247**(3-4): 382-401.
- 487 Murray, J.W. 2001. The niche of benthic foraminifera, critical thresholds and proxies.
488 *Marine Micropaleontology*, **41**: 1-7.
- 489 Murray, J.W. 2006. Ecology and Applications of Benthic Foraminifera. Cambridge
490 University Press, Cambridge (UK).
- 491 Nees, S. & Struck, U. 1994. The biostratigraphic and paleoceanographic significance of
492 *Siphotextularia rolshauseni* Phleger and Parker in Norwegian-Greenland Sea sediments.
493 *Journal of Foraminiferal Research*, **24**(4): 233-240.
- 494 Nees, S. 1997. Late Quaternary palaeoceanography of the Tasman Sea: The benthic
495 foraminiferal view. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **131**(3-4): 365-389.
- 496 Paillard, D., Labeyrie, L. & Yiou, P. 1996. Macintosh Program Performs Time-Series
497 Analysis. Eos, Transactions, American Geophysical Union, **77**: 379.
- 498 Parker, F.L. 1958. Eastern Mediterranean Foraminifera. *Reports of the Swedish Deep Sea*
499 *Expedition* **8**, 217-283.
- 500 Peterson, L.C. 1983. Recent Abyssal benthic foraminiferal biofacies of the Eastern
501 Equatorial Indian Ocean. *Marine Micropaleontology*, **8**: 479-519.
- 502 Phleger, F.B., Parker, F.L. & Peirson, J.F. 1953. North Atlantic foraminifera. *Reports of*
503 *the Swedish Deep Sea Expedition*, **7**:1-122.

- 504 Poli, M.S., Meyers, P.A., Thunell, R.C. & Capodivacca, M. 2012. Glacial-interglacial
505 variations in sediment organic carbon accumulation and benthic foraminiferal assemblages on
506 the Bermuda Rise (ODP Site 1063) during MIS 13 to 10. *Paleoceanography*, **27**(3): PA3216,
507 doi: 10.1029/2012pa002314.
- 508 Rasmussen, T.L., Oppo, D.W., Thomsen, E. & Lehman, S.J. 2003. Deep sea records from
509 the southeast Labrador Sea: Ocean circulation changes and ice-rafting events during the last
510 160, 000 years. *Paleoceanography*, **18**(1): 18-11.
- 511 Rathburn, A.E. & Corliss, B.H. 1994. The ecology of living (stained) deep-sea benthic
512 foraminifera from the Sulu Sea. *Paleoceanography*, **9**: 87-150.
- 513 Sachs, J.P. & Anderson, R.F. 2003. Fidelity of alkenone paleotemperatures in southern
514 Cape Basin sediment drifts. *Paleoceanography*, **18**(4): 1082, doi: 10.1029/2002pa000862.
- 515 Sachs, J.P. & Anderson, R.F. 2005. Increased productivity in the subantarctic ocean
516 during Heinrich events. *Nature*, **434**(7037): 1118-1121.
- 517 Schmiedl, G. & Mackensen, A. 1997. Late Quaternary paleoproductivity and deep water
518 circulation in the eastern South Atlantic Ocean: Evidence from benthic foraminifera.
519 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **130**: 43-80.
- 520 Schmiedl, G., Mackensen, A. & Müller, P.J. 1997. Recent benthic foraminifera from the
521 eastern South Atlantic Ocean: Dependence on food supply and water masses. *Marine*
522 *Micropaleontology*, **32**(3-4): 249-287.
- 523 Schönfeld, J., Alve, E., Geslin, E., Jorissen, F., Korsun, S. & Spezzaferri, S. 2012. The
524 FOBIMO (FORaminiferal BIo-MONitoring) initiative—Towards a standardised protocol for
525 soft-bottom benthic foraminiferal monitoring studies. *Marine Micropaleontology*, **94–95**: 1-
526 13.
- 527 Smart, C.W., King, S.C., Gooday, A.J., Murray, J.W. & Thomas, E. 1994. A benthic
528 foraminiferal proxy of pulsed organic matter paleofluxes. *Marine Micropaleontology*, **23**: 89-
529 99.
- 530 Smart, C.W. & Gooday, A.J. 1997. Recent benthic foraminifera in the abyssal Northeast
531 Atlantic Ocean; relation to phytodetrital inputs. *Journal of Foraminiferal Research*, **27**(2): 85-
532 92.
- 533 Smart, C.W. 2008. Abyssal NE Atlantic benthic foraminifera during the last 15 kyr:
534 Relation to variations in seasonality of productivity. *Marine Micropaleontology*, **69**: 193-211.

- Smart, C.W., Waelbroeck, C., Michel, E. & Mazaud, A. 2010. Benthic foraminiferal abundance and stable isotope changes in the Indian Ocean sector of the Southern Ocean during the last 20kyr: Paleoceanographic implications. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **297**(3–4): 537-548.
- Smith, P.B. 1964. Ecology of Benthonic species : A discussion of depth distribution of Foraminifera and ecologic factors off El Salvador, including a comparison with other Pacific coast areas. U.S. Government Printing Office.
- Smith, C.R., De Leo, F.C., Bernardino, A.F., Sweetman, A.K. & Arbizu, P.M. 2008. Abyssal food limitation, ecosystem structure and climate change. *Trends in ecology & evolution*, **23**(9): 518-528.
- Struck, U. 1995. Stepwise postglacial migration of benthic foraminifera into the abyssal northeastern Norwegian Sea. *Marine Micropaleontology*, **26**(1–4): 207-213.
- Suhr, S.B., Pond, D.W., Gooday, A.J. & Smith, G.R. 2003. Selective feeding by benthic foraminifera on phytodetritus on the western Antarctic Peninsula shelf: evidence from fatty acid biomarker analysis. *Marine Ecology Progress Series*, **262**: 153-162.
- Sun, X., Corliss, B.H., Brown, C.W. & Showers, W.J. 2006. The effect of primary productivity and seasonality of deep-sea benthic foraminifera in the North Atlantic. *Deep-Sea Research I*, **53**: 28-47.
- Szarek, R., Kuhnt, W., Kawamura, H. & Nishi, H. 2006. Distribution of recent benthic foraminifera along continental slope of the Sunda Shelf (South China Sea). *Marine Micropaleontology*, **71**(1–2): 41-59.
- Thiel, H., Pfannkuche, O., Schriever, G., Lochte, K., Gooday, A.J., Hemleben, C., Mantoura, R.F.G., Turley, C.M., Patching, J.W. & Riemann, F. 1989. Phytodetritus on the Deep-Sea Floor in a Central Oceanic Region of the Northeast Atlantic. *Biological Oceanography*, **6**(2): 203-239.
- Thomas, E., Booth, L., Maslin, M. & Shackleton, N.J. 1995. Northeastern Atlantic benthic foraminifera during the last 45,000 years: Changes in productivity seen from the bottom up. *Palaeoceanography*, **10**(3): 545-562.
- Zarriess, M. & Mackensen, A. 2010. The tropical rainbelt and productivity changes off northwest Africa: A 31,000-year high-resolution record. *Marine Micropaleontology*, **76**(3–4): 76-91.

FIGURE CAPTIONS:

Figure 1:

Location of core TN057-21 in the South Atlantic Southern Ocean (map elaborated using Ocean Data View, Schlitzer, R., <http://odv.awi.de>, 2015). Detailed information about modern hydrography and primary productivity patterns can be found in Llido *et al.* (2004) and Machu *et al.* (2005).

Figure 2: Benthic foraminiferal assemblages in core TN057-21

Shannon-H diversity index (a) and the relative abundance of *Epistominella exigua* (b) and the most characteristic species of benthic foraminifera in core TNO57-21 (c-n). Percentages are shown in the total assemblage (grey line) and in the phytodetritus free assemblages (black filled area). Percentages are calculated on the basis of samples containing > 50 specimens and non affected by dissolution. Intensely dissolved intervals that correspond to Greenland Stadials 19 and 21 are indicated by vertical bars. The ecological preferences for the most characteristic benthic foraminiferal species or group of species are summarized in Table 1.

Figure 3: Paleoproductivity from benthic foraminiferal accumulation rates and assemblage composition

The Benthic Foraminiferal Accumulation Rate (BFAR, total number of foraminifera >125 $\mu\text{m cm}^{-2} \text{ kyr}^{-1}$) is indicated as a shaded black areas and Benthic Foraminiferal Accumulation Rate calculated on phytodetritus free basis (i.e., discounting the contribution of the phytodetritus-

related species *Epistominella exigua*, $\text{BFAR}_{\text{phfree}}$) is indicated by a grey shaded area (a). Consequently, the difference between the two shaded areas is the BFAR of *E. exigua* ($\text{BFAR}_{E.exigua}$). Figure c displays the flux of alkenones ($\text{ng cm}^{-2} \text{ kyr}^{-1}$) in core TNO57-21, a proxy for organic carbon flux to the seafloor (Sachs & Anderson, 2005) plotted on Barker & Diz (2014) age model. The group of "low flux" species (e) is composed by shallow infaunal *O. umbonatus*, *G. subglobosa* and epifaunal *P. murrinha* and *F. wuellerstorfi*. Intermediate flux species (light grey line, d) is constituted by individuals belonging to *P. osloensis*, *S. rolhsauseni* (medium grey line) and *Melonis* spp. (dark grey line). Percentages are calculated on the phytodetritus free assemblage basis. Records of Southern Hemisphere ice core temperature (f, Antarctica Epica Dome C δD , Jouzel *et al.*, 2007, 3 points running average) and Northern Hemisphere ice core temperature record (b, Greenland NGRIP $\delta^{18}\text{O}$; EPICA, 2006) are showed as reference. Figures b and f are plotted on GICC05/NALPS timescale (Barker *et al.*, 2011). Black crosses on the top indicate tie points and sedimentation rates respectively used in the age model construction and BFAR calculations. Grey line on the top plot shows density values calculated using model proposed by Sachs & Anderson (2003).

TABLE CAPTIONS:

Table 1: Ecological attributions of the most characteristic benthic foraminiferal species in core TNO57-21..


	SPECIES	MICROHABITAT	ECOLOGICAL SIGNIFICANCE
Low Flux	<i>Fontbotia wuellerstorfi</i> (Schwager)= <i>Anomalina wuellerstorfi</i> Schwager, 1866. See Phleger <i>et al.</i> 1953, Pl. 11, Fig. 1-2. This work, Plate I, Fig. 8.	Epifaunal/ Epibenthic	<i>F. wuellerstorfi</i> is an epibenthic species (Linke & Lutze, 1993) that lives in well oxygenated and ventilated bottom waters away from high productivity coastal environments. Global distributional patterns indicate that it prefers organic fluxes below 2 gCm ⁻² yr ⁻¹ (Altenbach <i>et al.</i> 1999). In the South Atlantic, the distribution of this species coincides with relatively sustained food fluxes in areas where carbonate dissolution is not too severe (Mackensen <i>et al.</i> , 1995). In deep sea sediments of the Indian Ocean <i>F. wuellerstorfi</i> co-occurs with the phytodetritus species <i>E. exigua</i> . On ocean wide scale it co-occurs with <i>Uvigerina peregrina</i> (a typical high-flux species) on the lower range of organic carbon fluxes of the later (Altenbach <i>et al.</i> , 1999). This information suggest that <i>F. wuellerstorfi</i> is adapted to a wide range of environmental conditions, including strongly pulsed supply of organic carbon in the Indian Ocean (Gupta, 1997; De & Gupta, 2010).
Low Flux	<i>Oridorsalis umbonatus</i> (Reuss)= <i>Rotalina umbonata</i> Reuss 1851. See Smith, 1964, Pl. 4, Fig. 8 as <i>Pseudoepionides umbonatus</i> . This work, Plate I, Fig. 15. Most of the individuals found in core TN057-21 are juvenile forms.	Shallow infaunal	On an ocean wide scale, <i>O. umbonatus</i> seems to be distributed in areas receiving low organic carbon fluxes (Altenbach <i>et al.</i> 1999). Mackensen <i>et al.</i> (1995) suggest that the distribution of <i>O. umbonatus</i> in the South Atlantic is related to relatively well-oxygenated pore waters receiving a relatively low but sustained food supply. It co-occurs with <i>E. exigua</i> in the carbonate supersaturated and low organic carbon areas of the Weddell sea continental Shelf (Mackensen <i>et al.</i> 1990). In the eastern Indian Ocean, Murgese & DeDecker (2005) found the assemblage composed by <i>O. umbonatus</i> - <i>E. exigua</i> and <i>P. murrhina</i> indicator of low organic carbon. In Sulu area, Miao & Thunell (1993) indicated <i>O. umbonatus</i> as a species inhabiting shallow infaunal low organic carbon, well oxygenated sediments in agreement with Rathburn & Corliss (1994). Likewise, Burke <i>et al.</i> (1993), include <i>O. umbonatus</i> together with <i>Eggerella bradyi</i> within the group of low productivity species in the central Pacific. Conversely Gupta (1997) interpret the group of <i>O. umbonatus</i> , <i>Eggerella bradyi</i> , and <i>Melonis pompilioides</i> as reflect intermediate flux of relatively degraded organic matter and Kaiho (1999) consider <i>O. umbonatus</i> as an indicator of low-oxygen conditions.
Low Flux	<i>Pyrgo murrhina</i> (Schwager)= <i>Biloculina murrhina</i> Schwager 1866. See Phleger <i>et al.</i> 1953, Pl. 5, Fig. 22-24.	Epifaunal	<i>Pyrgo murrinha</i> is an epifaunal species (Corliss & Chen, 1988) found in areas of low organic carbon fluxes (Altenbach <i>et al.</i> , 1999) and high oxygenation such as the South Central Indian Ocean (De & Gupta, 2010) or eastern Indian Ocean (Murgese & De Deckker, 2005). Conversely, Gupta & Thomas (2003) interpret the abundance of <i>P. murrinha</i> in Indian cores as related to pulsed food and good carbonate preservation.
Low Flux	<i>Globocassidulina subglobosa</i> (Brady)= <i>Cassidulina subglobosa</i> Brady, 1881. See Parker 1958, Pl. 4, Fig. 13. This work, Plate I, Fig. 10.	Epifaunal/ Infaunal	In the North Atlantic Sun <i>et al.</i> (2006) found an inverse correlation of the <i>G. subglobosa</i> - <i>N. umbonifera</i> assemblage with seasonality of the organic matter. In the Southeast Atlantic, this species characterize sandy sediments of the Walvis Ridge (South Atlantic) with low organic carbon and high calcium carbonate content and enhanced bottom current velocities (Schmiedl <i>et al.</i> 1997). However, <i>G. subglobosa</i> has been associated with deposition of phytodetritus in the Northeast Atlantic Abyssal Plain Gooday (1988, 1993), North West Africa (Eberwein & Mackensen (2006) and the Antarctic shelf (Suhr <i>et al.</i> 2003).
Episodic phytoplankton blooms	<i>Epistominella exigua</i> (Brady)= <i>Pulvinulina exigua</i> Brady, 1884. See Lobegeier & Sen Gupta, 2008, Pl. I, Fig. 18. This work, Plate I, Fig. 9.	Epifaunal/ Shallow infaunal	This is an opportunistic species that rapidly colonizes and feeds on aggregates of phytodetritus produced by episodic phytoplankton blooms (e.g., Gooday, 1988, 1993, Cornelius & Gooday, 2004; Sun <i>et al.</i> 2006; Gooday <i>et al.</i> 2010). <i>Epistominella exigua</i> is well adapted to oligotrophic conditions well away from areas where the productivity is high and the flux of organic matter is continuous (Mackensen <i>et al.</i> 1995). In the eastern South Atlantic <i>E. exigua</i> is found in association with <i>F. wuellerstorfi</i> in low organic carbon areas on the flanks of the Walvis Ridge (Schmiedl <i>et al.</i> 1997). It is also abundant in the deep western South Atlantic (Harloff & Mackensen, 1997) the deep Weddell Sea continental slope (Anderson, 1975; Cornelius & Gooday, 2004) and South West Indian Ocean (Peterson, 1983; Corliss, 1983).
Intermediate Flux	<i>Pullenia osloensis</i> Feyling-Hanssen, 1954. See Diz <i>et al.</i> 2007- Supplementary information, Pl. II, Fig. 5. This work, Plate I, Fig. 16.	Not very well constrained Shallow to deep infaunal	The ecology of <i>Pullenia osloensis</i> is not well known. This species is described in deep and abyssal areas of the South Cape Basin (Boltovskoy & Boltovskoy, 1989), Southwest Indian Ocean (Corliss, 1979) and the Agulhas retroflection area (Diz <i>et al.</i> 2007). Most authors interpret the group of <i>Pullenia</i> spp. (mainly <i>Pullenia bulloides</i>) as indicator of areas of high organic supply (Rasmussen <i>et al.</i> 2003) or high but varying surface productivity (Mackensen <i>et al.</i> 1995). In the Indian Ocean, the assemblage dominated by <i>Epistominella exigua</i> with <i>Pullenia osloensis</i> and <i>Pullenia salisburyi</i> as secondary species is considered indicator of low to intermediate organic flux and high seasonality (Gupta & Thomas, 2003). <i>Pullenia osloensis</i> is common along core TNO57-21 suggesting that it is adapted to a wide range of carbon flux regimes (from low to intermediate) and organic carbon qualities (i.e., labile and refractory). <i>Pullenia salisburyi</i> is also present in core TNO57-21 (Plate, Fig. 17) even though its relative contributions are lower than <i>P. osloensis</i> .
Intermediate Flux	<i>Melonis zaandamae</i> (Van Voothuysen) = <i>Melonis barleeaanum</i> var. <i>zaandamae</i> van Voothuysen, 1952. See Mackensen <i>et al.</i> 1993, Pl.3, Fig. 4-5. This work, Plate I, Fig. 13.	Shallow infaunal	<i>Melonis</i> spp. species (in this core mainly <i>M. zaandamae</i>) are intermediate infaunal foraminifera that tolerates moderate oxygen depletion (Fontanier <i>et al.</i> , 2002). It is generally associated to intermediate organic carbon flux and refractory organic matter (Caralp, 1989; Alve, 2010). Mackensen <i>et al.</i> (1993, 1995) refer to the assemblage comprising <i>Melonis pompilioides</i> , <i>M. zaandamae</i> , and <i>O. umbonatus</i> as South Atlantic deep sea high productivity fauna.
Intermediate Flux	<i>Siphotextularia rolshauseni</i> Phleger and Parker, 1951. See Phleger <i>et al.</i> 1953, Pl. 5, Fig. 7. This work, Plate I, Fig. 2.	Not well constrained	<i>Siphotextularia rolshauseni</i> is an agglutinated species composed of calcareous fragments of various sizes and it is considered a junior synonym of <i>Siphotextularia catenata</i> Cushman (Corliss, 1979, Nees & Struck, 1994). There are a few references to this species in superficial sediments of the North Atlantic (see review Nees & Struck, 1994), Southwest Pacific (Kurihara & Kennett, 1986), South China Sea (Szarek <i>et al.</i> , 2006), the Indian Ocean (Corliss, 1979, Nees <i>et al.</i> 1997). Similarly, this species is reported as rare in South Atlantic superficial sediments by Mackensen <i>et al.</i> (1993, <i>Siphotextularia</i> sp.), Schmiedl <i>et al.</i> (1997, <i>S. catenata</i>) as well as in Pleistocene and Miocene sediments of the Walvis Ridge (Boltovskoy & Boltovskoy, 1989). Nees & Struck (1994) and Struck (1995) relate the occurrence of <i>S. rolshauseni</i> in the Greenland-Norwegian Seas during MIS 2 with a period of low trophic quality. This interpretation is also suggested by the downcore distribution of <i>S. rolshauseni</i> in core TNO57-21 which peaks together with <i>Melonis</i> spp. during MIS 4 (see text).
Flux Range?	The called "Lagenina Group" is composed by several species belonging to genus <i>Amphycorina</i> , <i>Bifarilaminella</i> , <i>Cushmanina</i> , <i>Fissurina</i> , <i>Homaloedra</i> ;	Unknown	The species belonging to the called "Lagenina group" are found as rare representatives of the living and recently dead assemblages in different marine settings. Thus, the ecological significance of this group is not well constrained. Some of the species belonging to Lagenina group are illustrated in Plate I: <i>Fissurina staphyllearia</i> (Plate, Fig. 3),


615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636

PLATE I CAPTION:

PLATE I:

1. *Eggerella bradyi* (Cushman), scale bar 100 μm .
2. *Siphotextularia rolshauseni* Phleger and Parker, scale bar 100 μm .
3. *Fissurina staphyllearia* Schwager, scale bar 100 μm .
4. *Oolina globosa* (Montagu), scale bar 100 μm .
5. *Lagena hispidula* Cushman, 1913, scale bar 100 μm .
6. *Parafissurina felsinea* (Fornasini), scale bar 100 μm .
7. *Quinqueloculina* cf. *seminula*, scale bar 100 μm .
8. *Fontbotia wuellerstorfi* (Schwager), scale bar 100 μm .
9. *Epistominella exigua* (Brady), 9a, spiral side, 9b umbilical side, scale bars 100 μm .
10. *Globocassidulina subglobosa* (Brady), scale bar 200 μm .
11. *Rutherfordoides tenuis* (Phleger and Parker), scale bar 200 μm .
12. *Gyroidinoides* cf. *polius*, scale bar 100 μm .
13. *Melonis zaandamae* (Van Voorthuysen), scale bar 100 μm .
14. *Nutallides umboniferus* (Cushman), scale bar 100 μm .
15. *Oridorsalis umbonatus* (Reuss), 4a, umbilical side, 4b spiral side, scale bars 100 μm .
16. *Pullenia osloensis* Feyling-Hanssen, scale bar 100 μm .
17. *Pullenia salisburyi* Stewart and Stewart, scale bar 100 μm .
18. *Uvigerina auberiana* d'Orbigny, scale bar 200 μm .








Plate I

